Preventing PICS: Targeted QI in the ICU

Nida Qadir, MD Assistant Professor of Medicine Associate Director, Medical ICU Co-Director, Post-ICU Recovery Clinic Ronald Reagan UCLA Medical Center





1

ABCDEF bundle

- Assess, prevent, and manage pain
- Both spontaneous awakening and breathing trials
- Choice of analgesia and sedation
- Delirium: assess, prevent, and manage
- Early mobility and exercise
- Family empowerment and engagement

Ely, Crit Care Med 2017





ICU Liberation Collaborative

- 15,226 patients
- 68 ICUs
- Complete bundle performance → improved outcomes

Pun et al,	Crit Care	<i>Med</i> 2019
------------	-----------	-----------------





Complete Bundle Performance

AHR (95% CI)

1.17 (1.05–1.30)

1.19(1.01-1.40)

0.32 (0.17-0.62)

AOR (95%CI)

0.28 (0.22-0.36)

0.35 (0.22–0.56)

0.60 (0.49–0.72)

1.03 (0.88–1.21)

0.37 (0.30–0.46)

0.54 (037-0.79)

0.64 (0.51-0.80)

Adjusted OR (95%CI)

Outcomes

Death^c

Coma

Delirium

Significant pain

Physical restraints

ICU readmission^e

System-Related Outcomes

Discharge destination¹

Patient-Related Outcomes

ICU discharge^a

Hospital discharge^b

Symptom-Related Outcomes^d

Mechanical ventilation

p Value

< 0.004

< 0.033

< 0.001

< 0.0001

< 0.0001

< 0.0001


< 0.0001

< 0.001

< 0.001

0.7000

ICU Liberation Collaborative



Pun et al, Crit Care Med 2019





Ensure smooth transitions in care

Medications

- Resume chronic meds
- Stop unnecessary meds
- Adjust medication doses
- PT/OT assessment
- Tubes/lines/devices
- Identify caregiver
- Arrange follow-up
- Educate about PICS

Bell et al, *JAMA*Scales et al *J Gen Intern Med*Taylor et al *Ann Am Thorac Soc*Kovacic *J Pharm Pract*





How post-ICU activities can improve ICU care

- Identifying otherwise unseen targets for ICU QI
- Creating a new role for survivors in the ICU
- Inviting ICU providers to the post-ICU program as a learning strategy
- Challenging clinician's own understanding of patient experience
- Improving morale and meaningfulness of ICU work

Haines et al, Intensive Care Med 2019





Feedback from PICS clinic

- 53M with h/o HTN admitted to the ICU with COVID-19 PNA and severe ARDS
- Intubated x 10 days
- Discharged to rehab facility
- Seen post-ICU clinic 6 weeks later





Feedback \rightarrow QI

- Neuropathy in R wrist (site of arterial line)
 - Updated inpatient ICU guidance to recommend arterial line placement on non-dominant side
- Facial pressure ulcers from proning
 - Discussed with inpatient RN team \rightarrow proning protocol modified
- Returned home when many thought his illness was not survivable
 - Improves GOC discussions
 - Improves morale!





Thank you!



